

INSTRUCTIONS ON USE OF

DERMISIL® R TOPICAL RINGWORM TREATMENT

Always Patch Test an Area Before Starting Full Treatment Regiment

Wash affected area with soap and water. Washing the area every time is an important part of the treatment regiment. Dry affected area completely. Keeping the area dry is a very important part of successfully removing the ringworm fungus. Wash the area with hydrogen peroxide solution to ensure a very effective treatment program and that the area is clean prior to treatment.

Be sure to wash any bedding, clothes, laundry etc during the treatment time because spores can easily reinfect you or your family members without proper hygiene. Animals also carry the ringworm fungal spores and may need to be treated as well.

BenzaRid® is a anti-fungal spray that kills fungal spores on contact. It can be spray on carpet, bedding, and anywhere you suspect fugal spores to live.

Apply Dermisil® R liquid to the affected areas with a clean cotton ball, or q-tip.

Do not use the Dermisil R product on animals. The Dermisil R for Pets is the only product that should be used on animals.

Apply 3 times or more per day until the ringworm fungus is destroyed. The fungus should start to diminish within 48-72 hours depending on the severity of the outbreak. It may take up to several weeks to completely eliminate the ringworm.

You may have drying of the skin area where the liquid is applied. This is normal and will subside after the treatment regiment is completed.

The damage left behind from ringworm will slowly dissipate over several weeks, depending on the level of outbreak. If hair follicles have been destroyed by the ringworm fungus the hair should return within several weeks to months. There may be scarring damage left from the fungus, unfortunately this is normal depending on the severity and length of time the ringworm was present.

If for some reason the area has not diminished in size over several weeks, you may need to seek a dermatologist for a culture of the area. You may not have ringworm, but another skin disorder. We can not be responsible for a mis-diagnosis of your condition.

Note: Active ingredients in DERMISIL® R are: cymbopogon citratus, melaleuca alternifolia, and other anti-fungal plant extracts .

NEVER apply Dermisil® R liquid to children under 5 years of age or if you are pregnant or nursing. NEVER apply Dermisil® R liquid to mucous membranes or to the eyes. NEVER ingest or drink Dermisil® R liquid. If Dermisil® R is ingested immediately phone the Poison Control Center.