

INSTRUCTIONS ON USE OF DERMISIL® W TOPICAL WART TREATMENT FOR COMMON WARTS, GENITAL WARTS, FLAT WARTS, SKIN TAGS, MOLLUSCUM

FOR WARTS

Always Patch Test an Area Before Starting Full Treatment Regimen

Wash affected area with soap and water for at least several minutes.

Dry affected area completely.
Use an emery board to score the wart a little. This will remove some of the dead tissue prior to treatment.

Perform the same emery board treatment in the morning and evening, with a topical treatment during the day. If you skip a treatment, it could set your recovery back significantly.

Apply Dermisil® W liquid to the affected areas with a q-tip 4 times per day. **At night, cut a small piece of cotton ball the size of the wart and dip it in the Dermisil® W liquid, then use a piece of medical tape to secure it to the wart area, if possible. This will speed the process greatly.** The wart may increase in size for a short period of time while the virus fights the treatment. This is normal. The wart should start to diminish within 2 to 6 weeks depending on the severity of your wart condition. Remember, there are over 100 different types of the wart virus and HPV that causes warts. Some wart types can be quite virulent. Keep applying the treatment until the wart is completely gone.

You may have drying of the skin area where the liquid is applied as the wart flakes away. This is normal and will subside after the treatment regimen is completed.

An Immune System Booster such as Dermisil Immune Renew can speed the process of wart removal which will help your body fight the HPV virus.

The damage left behind from the wart will diminish over time, depending on the level of outbreak.

Do not apply to the eye area.

FOR PLANTAR'S WARTS

Soak and wash affected area with soap and water for at least several minutes. Dry affected area completely. Use an emery board to score the wart every day. This will remove some of the dead tissue prior to daily/nightly treatments.

Apply Dermisil® W liquid to the affected areas with a clean cotton ball, or q-tip 4 times per day. **At night, cut a small piece of cotton ball the size of the wart and dip it in the Dermisil® W liquid, then use a piece of medical tape to secure it to the bottom of the foot every night at bed-time. This will speed the process greatly.** There may be some burning sensation, but do not be alarmed.

If you skip a treatment, it could set your recovery back significantly.

Perform the same emery board treatment in the morning and evening, with a topical treatment during the day. Apply 4 times or more per day until the wart is destroyed. The wart should start to diminish within 2 to 6 weeks depending on the severity of your wart condition. The wart may increase in size for a short period of time while the virus fights the treatment. This is normal. You may have drying of the skin area where the liquid is applied as the wart flakes away. This is normal and will subside after the treatment regimen is completed.

The damage left behind from the wart will diminish over time, depending on the level of outbreak.

FOR SKIN TAGS AND MOLLUSCUM

Wash affected area with soap and water for at least several minutes. Dry affected area completely.

Apply Dermisil® W liquid to the affected areas with a clean cotton ball, or q-tip. There may be some burning sensation, but do not be alarmed.

If you skip a treatment, it could set your recovery back significantly. Apply 4 times or more per day **At night, cut a small piece of cotton ball the size of the skin tag or molluscum and dip it in the Dermisil® W liquid, then use a piece of medical tape to secure it to the area, if possible. This will speed the process greatly.** Keep applying the treatment until the skin-tag or molluscum is completely gone. You may have drying of the skin area where the liquid is applied as the wart flakes away. This is normal and will subside after the treatment regimen is completed. The damage left behind from the skin-tag or molluscum will diminish over time, depending on the level of outbreak. **Do not apply to the eye area.**

FOR GENITAL WARTS

Always Patch Test an Area Before Starting Full Treatment Regimen

Wash affected area with soap and water for at least several minutes.

Dry affected area completely. Apply Dermisil® W liquid to the affected genital areas with a q-tip. There may be some burning sensation, but do not be alarmed. If you skip a treatment, it could set your recovery back significantly.

If you have internal genital warts, anal, or vaginal warts, you may try using a tampon with a small amount of liquid on the tampon and apply it to wart area. Always test and area first before applying fully. It is possible that you may need to seek professional medical attention for the removal of internal genital warts.

Apply 4 times or more per day until the warts are destroyed. The warts should start to diminish within 2 to 6 weeks depending on the severity of your wart condition.

The warts may increase in size for a short period of time while the virus fights the treatment. This is normal.

Remember, there are over 100 different types of the wart virus and HPV that causes warts. Some wart types can be quite virulent. Keep applying the treatment until the warts are completely gone.

You may have drying of the skin area where the liquid is applied as the wart flakes away. This is normal and will subside after the treatment regimen is completed.

The damage left behind from the wart will diminish over time, depending on the level of outbreak. **An Immune System Booster such as Dermisil Immune Renew can speed the process of wart removal which will help your body fight the HPV virus.**

Note: Active ingredients in DERMISIL® W are: thuja occidentalis, melaleuca alternifolia, and ricinus communis plant extracts. NEVER apply Dermisil® W liquid to children under 5 years of age or if you are pregnant or nursing. NEVER apply Dermisil® W liquid to the eyes or eye area. NEVER ingest or drink Dermisil® W liquid. If Dermisil® W is ingested immediately phone the Poison Control Center.